

An easier solution of a Diophantine problem about triangles, in which those lines from the vertices which bisect the opposite sides may be expressed rationally*

Leonhard Euler[†]

1. The sides of a triangle having been placed as $AB = 2c$, $AC = 2b$ and $BC = 2a$, if the bisecting lines are called $AX = x$, $BY = y$ and $CZ = z$, it is apparent from Geometry for the squares of these three lines to be expressed in the following way:

$$xx = 2bb + 2cc - aa$$

$$yy = 2cc + 2aa - bb$$

$$zz = 2aa + 2bb - cc,$$

and these equations are thus to be resolved through rational numbers.

*Delivered to the St.-Petersburg Academy on August 12, 1779. Originally published as *Solutio facilior problematis Diophantei circa triangulum, in quo rectae ex angulis latera opposita bisecantes rationaliter exprimantur*, Mémoires de l'Académie Impériale des Sciences de St.-Pétersbourg **2** (1810), 10-16, and reprinted in *Leonhard Euler, Opera Omnia*, Series 1: Opera mathematica, Volume 4, Birkhäuser, 1992. A copy of the original text is available electronically at the Euler Archive, at www.eulerarchive.org. This paper is E732 in the Eneström index.

[†]Date of translation: March 2, 2005. Translated from the Latin by Jordan Bell, 2nd year undergraduate in Honours Mathematics, School of Mathematics and Statistics, Carleton University, Ottawa, Ontario, Canada. Email: jb3@connect.carleton.ca. Translation written under Dr. B. Mortimer.

2. From these three equations, the following three are formed:

$$\begin{aligned} \text{I. } & xx - yy = 3(bb - aa) \\ \text{II. } & xx + yy = 4cc + aa + bb \\ \text{III. } & zz = 2aa + 2bb - cc. \end{aligned}$$

We deal with these equations in the following way.

3. We start with the first of these equations, and so that we can avoid fractions, we set $xx - yy = 3(bb - aa) = 12fg(pp - qq)$; from this, with it $bb - aa = 4fg(pp - qq)$, we make it $b + a = 2f(p + q)$ and $b - a = 2g(p - q)$, from which it will be $b = (f + g)p + (f - g)q$ and $a = (f - g)p + (f + g)q$. Indeed, with the previous equation $xx - yy = 12fg(pp - qq)$, we take $x + y = 6g(p + q)$ and $x - y = 2f(p - q)$, from which it will be $x = (3g + f)p + (3g - f)q$ and $y = (3g - f)p + (3g + f)q$.

4. The second equation is now approached:

$$xx + yy = 4cc + aa + bb,$$

and together with the values that have just been determined, it is discovered that:

$$xx + yy = 2(9gg + ff)(pp + qq) + 4(9gg - ff)pq.$$

Then indeed it will be:

$$bb + aa = 2(ff + gg)(pp + qq) + 4(ff - gg)pq.$$

Therefore, with it $4cc = xx + yy - (bb + aa)$, it will be:

$$4cc = 16gg(pp + qq) + (40gg - 8ff)pq,$$

on account of which we have $cc = 4gg(pp + qq) + (10gg - 2ff)pq$, so that therefore this formula is to be transformed into a square.

5. From here the third equation is proceeded to, which is $zz = 2(aa + bb) - cc$, from which it will be:

$$\begin{aligned} 2(aa + bb) &= 4(ff + gg)(pp + qq) + 8(ff - gg)pq \quad \text{and} \\ cc &= 4gg(pp + qq) + (10gg - 2ff)pq, \quad \text{and it will be} \\ zz &= 4ff(pp + qq) + (10ff - 18gg)pq, \end{aligned}$$

which is thus another formula that we must reduce to a square.

6. We divide the former of these two equations by $4gg$ and the latter indeed by $4ff$, so that we may have the following formulas, reduced into squares:

$$\begin{aligned}\frac{cc}{4gg} &= pp + qq + 2pq \left(\frac{5gg - ff}{4gg} \right) \\ \frac{zz}{4ff} &= pp + qq + 2pq \left(\frac{5ff - 9gg}{4ff} \right).\end{aligned}$$

7. For briefness, here we put $\frac{5gg - ff}{4gg} = m$ and $\frac{5ff - 9gg}{4ff} = n$, so that now the entire problem is brought to the resolution of these two formulas:

$$\begin{aligned}\frac{cc}{4gg} &= pp + qq + 2mpq = tt \\ \frac{zz}{4ff} &= pp + qq + 2npq = uu,\end{aligned}$$

and with this we will have $c = 2gt$ and $z = 2fu$. At this point, in order for these two formulas to be reduced to squares, it is noted to be that $tt - uu = 2(m - n)pq$, and for convenience it is thus permitted to treat this such that it is set $t + u = (m - n)p$ and $t - u = 2q$, from which is gathered $t = \frac{1}{2}(m - n)p + q$ and $u = \frac{1}{2}(m - n)p - q$.

8. But if these values are now substituted in place of t and u , it comes about that squares simultaneously arise on both sides of the equation:

$$pp \left(1 - \frac{1}{4}(m - n)^2 \right) + (m + n)pq = 0,$$

and when this equation is divided by p it is:

$$p \left(1 - \frac{1}{4}(m - n)^2 \right) + (m + n)q = 0,$$

from which the ratio between the letters p and q freely comes forth: $\frac{p}{q} = \frac{4(m+n)}{(m-n)^2 - 4}$, on account of which it will be possible for it to be taken $p = 4(m+n)$ and $q = (m - n)^2 - 4$, or equivalent multiples, so that it is to be held in general that $p = 4(m + n)M$ and $q = ((m - n)^2 - 4)N$, and in this way all the conditions are fully satisfied.

9. Now with the letters p and q that have been worked out, it will be $t = 2(m+n)(m-n) + (m-n)^2 - 4 = (m-n)(3m+n) - 4$ and $u = (m-n)(m+3n) + 4$, from which as before we infer these determinations:

$$\begin{aligned} c &= 2g(m-n)(3m+n) - 8g \quad \text{and} \\ z &= 2f(m-n)(m+3n) + 8f. \end{aligned}$$

Moreover, it is helpful to sum in the p and q places the values of them from §7, so that it will thus be $c = g(m-n)p + 2gq$ and $z = f(m-n)p - 2fq$.

10. Now therefore a solution of our problem is able to be constructed in the following way:

1.) Both the letters f and g are allowed to be at our choice, from which the letters m and n may be defined by means of the formulas $m = \frac{5gg-fg}{4ff}$ and $n = \frac{5ff-9gg}{4ff}$.

2.) Now as before the letters p and q are sought through the power of the formulas $p = 4(m+n)$ and $q = (m-n)^2 - 4$, with which having been obtained, the lines bisecting the sides of the triangle can be expressed in the following way.

3.) Of course the sides are found with these formulas:

$$\begin{aligned} a &= (f-g)p + (f+g)q \\ b &= (f+g)p + (f-g)q \\ c &= 2g(m-n)(3m+n) - 8g = g(m-n)p + 2gq \end{aligned}$$

4.) Finally therefore, the bisecting lines themselves will be had:

$$\begin{aligned} x &= (3g+f)p + (3g-f)q \\ y &= (3g-f)p + (3g+f)q \\ z &= 2f(m-n)(m+3n) + 8f = f(m-n)p - 2fg. \end{aligned}$$

11. So that we can illustrate an example of this, we suppose that $f = 2$ and $g = 1$, and it will then be $m = \frac{1}{4}$ and $n = \frac{11}{16}$. From this as before it is gathered that $p = \frac{15}{4}$ and $q = -\frac{975}{256}$, whose values, to be reduced to integral

numbers, are given as $p = 64$ and $q = -65$. Then our triangle will now be determined in this way:

$$\begin{aligned} a &= 131; b = 127; c = 158; \quad \text{then indeed,} \\ x &= 255; y = 261; z = 204. \end{aligned}$$

12. On this occasion as has been noted for the example, it will be helpful now too for the letters x, y, z to be taken for the sides, with in general it being:

$$\begin{aligned} 2xx - 2yy - zz &= 9cc \\ 2yy + 2zz - xx &= 9aa \\ 2zz + 2xx - yy &= 9bb, \end{aligned}$$

from which it follows, with it permitted for the numbers yielding x, y, z to be reduced by 3, then a very simple triangle is able to be formed, of which the sides are 136, 170, 174.

13. At this point, with it $m = \frac{5gg-ff}{4gg}$ and $n = \frac{5ff-9gg}{4ff}$, it will first be $m + n = \frac{10ffgg-f^4-9g^4}{4ffgg} = -\frac{(gg-ff)(9gg-ff)}{4ffgg}$. Then indeed it will be had that $m - n = \frac{9g^4-f^4}{4ffgg} = \frac{(3gg+ff)(3gg-ff)}{4ffgg}$, from which it is gathered that $m - n + 2 = \frac{9g^4+8ffgg-f^4}{4ffgg} = \frac{(gg+ff)(9gg-ff)}{4ffgg}$ and $m - n - 2 = \frac{9g^4-8ffgg-f^4}{4ffgg} = \frac{(gg-ff)(9gg+ff)}{4ffgg}$. By means of this, with it having been found that $p = 4(m+n)M$, it will now be:

$$\begin{aligned} p &= -\frac{(gg-ff)(9gg-ff)M}{4ffgg} \quad \text{and,} \\ q &= ((m-n)^2 - 4)M = \frac{(g^4-f^4)(81g^4-f^4)}{16f^4g^4}M. \end{aligned}$$

14. Therefore we now reduce the ratio of the letters p and q to minimum terms, and we take $M = \frac{16f^4g^4}{(gg-ff)(9gg-ff)}$, and thus it comes forth that $p = -16ffgg$ and $q = (gg+ff)(9gg+ff)$. From this, because it was $t = \frac{1}{2}(m-n)p + q$, it now follows that,

$$t = (gg+ff)(9gg+ff) - 2(3gg+ff)(3gg-ff).$$

In a similar way it will be $u = \frac{1}{2}(m-n)p - q$, from which it is gathered that:

$$u = -(gg + ff)(9gg + ff) - 2(3gg + ff)(3gg - ff).$$

Then because of this it will be $c = 2gt$ and $z = 2fu$.

15. For the remaining letters, it will first be $a + b = 2f(p + q)$ and $b - a = 2g(p - q)$; then indeed it will be $x + y = 6g(p + q)$ and $x - y = 2f(p - q)$. It is no problem for however many examples to be resolved readily enough by means of these formulas, to such a degree as they are according to integers; since they are in every respect determined by the ratio of the integral numbers f and g , these are therefore always allowed to be assumed integers.

16. It will be $f = 1$ and $g = 2$, and it will be $p = -64$ and $q = 185$. Hence now it is gathered $t = -101$ and $u = -471$; then indeed $c = -404$ and $z = -942$. To be sure, as before it will be $b - a = -996$ and $b + a = +242$; $x + y = 1452$ and $x - y = -498$. It will therefore be:

$$\begin{aligned} a &= 619; b = 377; c = 404 \\ x &= 477; y = 975; z = 942. \end{aligned}$$

It is observed that the numbers x, y, z are able moreover to be used instead of a, b, c , with these divided by 3, so that it becomes:

$$\begin{aligned} a &= 159; b = 325; c = 314, & \text{then also it will be,} \\ x &= 619; y = 377; z = 404. \end{aligned}$$